

The Theta-Gamma Neural Code

John E. Lisman^{1,*} and Ole Jensen^{2,*}

¹Brandeis University, Department of Biology & Volen Center for Complex Systems, 415 South Street-MS 008, Waltham, MA 02454-9110, USA

²Radboud University Nijmegen, Donders Institute for Brain, Cognition and Behaviour, Kapittelweg 29, 6525 EN Nijmegen, The Netherlands

*Correspondence: lisman@brandeis.edu (J.E.L.), ole.jensen@donders.ru.nl (O.J.)

<http://dx.doi.org/10.1016/j.neuron.2013.03.007>

Theta and gamma frequency oscillations occur in the same brain regions and interact with each other, a process called cross-frequency coupling. Here, we review evidence for the following hypothesis: that the dual oscillations form a code for representing multiple items in an ordered way. This form of coding has been most clearly demonstrated in the hippocampus, where different spatial information is represented in different gamma subcycles of a theta cycle. Other experiments have tested the functional importance of oscillations and their coupling. These involve correlation of oscillatory properties with memory states, correlation with memory performance, and effects of disrupting oscillations on memory. Recent work suggests that this coding scheme coordinates communication between brain regions and is involved in sensory as well as memory processes.

Introduction

Multi-item messages must often be transmitted between brain regions. For instance, short-term memory may represent the last several events in the recent past; similarly, the sequence of events that constitute an episodic memory may be recalled from long-term memory. Handling such multi-item messages requires a neural code that specifies not only how items are represented, but also how different items are kept separate (e.g., the longer pauses that separate letters in the Morse code). Here, we evaluate the hypothesis that the neural code for multi-item messages is organized by brain oscillations. These oscillations can be observed in field potentials, a method of extracellular recording that provides a measure of average neural activity in a brain region (Buzsáki et al., 2012). Such recordings in rodents (Figure 1A) have shown that gamma frequency (~40 Hz) oscillations are nested within slow theta frequency (~7 Hz) oscillations (Belluscio et al., 2012; Bragin et al., 1995; Colgin et al., 2009; Soltesz and Deschênes, 1993). A large number of experiments have investigated the role of theta/gamma oscillations, largely using physiological methods in rodents. More recently, the study of these oscillations in humans has become a focus of cognitive neuroscience (Axmacher et al., 2010; Canolty et al., 2006; Demiralp et al., 2007; Llinás and Ribary, 1993; Maris et al., 2011; Mormann et al., 2005; Sauseng et al., 2009; Voytek et al., 2010).

The specific hypothesis that we will evaluate here is shown in Figure 1B (Lisman and Buzsáki, 2008; Lisman and Idiart, 1995). According to this coding scheme, the subset of cells that fire during a given gamma cycle (sometimes referred to as a cell assembly or an ensemble) form a spatial pattern that represents a given item. Formatting of multiple items is organized by theta/gamma oscillations as follows: largely non-overlapping assemblies are active in different gamma cycles, i.e., at different theta phases. Given that there are four to eight gamma cycles nested within a theta cycle, multiple items can be represented in a defined order. Here, we will first describe the evidence that jointly occurring theta and gamma oscillations can organize information in the way hypothesized in Figure 1B. We will then describe experiments that address the following questions: (1)

do the oscillations and their interaction vary with cognitive demands, and do these changes predict behavioral performance? (2) Does interfering with (or enhancing) the oscillations affect function? (3) Are the oscillations used to coordinate communication between brain regions? We then turn to an analysis of the mechanistic role of gamma oscillations in the context of the theta-gamma code. In the final section, we discuss outstanding issues, notably the relationship of alpha and theta frequency oscillations in cortex and the possibility that the theta-gamma code contributes not only to memory processes, but also to sensory processes.

Theta-Gamma Coding in the Hippocampus

The first indication that theta oscillations have a role in neural coding came from the study of rat CA1 hippocampal place cells. Such cells increase their firing rate when the rat is in a subregion of the environment called the place field; different cells have different place fields (O'Keefe and Dostrovsky, 1971). As the rat crosses the place field of a cell, there are generally five to ten theta cycles. On each successive cycle, firing tends to occur with earlier and earlier theta phase (Figure 2A), a phenomenon termed the phase precession (O'Keefe and Recce, 1993; Skaggs et al., 1996). These and related results (Lenck-Santini et al., 2008; Pastalkova et al., 2008) suggest that the hippocampus uses a code in which theta phase carries information. Further analysis showed that CA1 place cells fire at a preferred phase of the faster gamma oscillations (Figure 2B; Senior et al., 2008). Thus, during a given theta cycle, firing will tend to occur at a preferred theta phase and at a preferred gamma phase.

If a place cell fires at a particular phase during a theta cycle (i.e., in a particular gamma cycle), other place cells representing different information presumably fire at other theta phases, collectively forming a multipart message. The ability to record simultaneously from >100 cells (Wilson and McNaughton, 1993) has made it possible to directly observe such messages. As illustrated in Figure 3, during an individual theta cycle, different place cells fire in a temporal sequence. These sequences are called "sweeps" because the firing order




Figure 1. Neural Code Organized by Theta and Gamma Oscillations (A) Simultaneous extracellular (top) and intracellular (bottom) recordings from the hippocampus. Intracellular gamma is due to IPSPs, the amplitude of which is modulated by the phase of theta. From Penttonen et al. (1998), used with permission.

(B) Schematic of the theta-gamma code. The ovals at top represent states of the same network during two gamma cycles (active cells are black and constitute the ensemble that codes for a particular item). Different ensembles are active in different gamma cycles.

corresponds to the cells' place field centers along the track (Gupta et al., 2012; see also Dragoi and Buzsáki, 2006; Harris et al., 2003; Skaggs et al., 1996). Such data show directly that different cells representing different information (i.e., positions) fire at different theta phases. Given that firing is also modulated by gamma (Figure 2B), there can be little doubt that a theta-gamma code is used in the hippocampus to represent ordered multi-item messages.

An important property of sweeps is that they are time compressed. Whereas the rat might take ~300 ms to move between positions a and b, cells representing these positions fire ~30 ms apart during a sweep (Skaggs et al., 1996). Furthermore, this time-compressed readout is often predictive (Battaglia et al., 2004), providing a way of rapidly informing downstream networks of the sequence of upcoming places (see Figure 2 caption). The usefulness of such predictive readout is particularly evident for the sweeps that occur when a rat stops at the choice point of a familiar maze (Johnson and Redish, 2007). Under these conditions, one sweep may represent the sequence of positions down one arm of the maze (sweeps can turn corners and thus are dependent on memory rather than direct vision), and the next sweep may be down the other arm. Such rapid readout from memory presumably allows downstream brain regions to choose the arm leading to the goal.

To be useful for transmitting information, a code must be consistent over cell populations and stable over time. A procedure for examining these requirements utilizes one part of the recording period to correlate potential coding variables (such as firing rate or theta phase) to the observed position of the rat. Different codes can then be quantitatively compared by their ability to predict the rat's position from the firing patterns during the other part of the experiment (Harris et al., 2003; Jensen and Lisman, 2000). It was found that codes that take theta phase into account allowed the rat's position to be predicted with an accuracy of about 3 cm, while codes that did not use theta phase had less accuracy. Together, these and related results (Harris et al., 2003) show that the theta-phase code carries information, is stable over time, and is used consistently by cell populations.




Figure 2. Spiking in the Rat CA1 Region Depends on the Phase of Both Theta and Gamma Oscillations

(A) Phase precession: as the rat runs through a place field, a process that takes several seconds, theta phase becomes progressively earlier (closer to zero phase) on successive theta cycles. From Mehta et al. (2002). The phase precession can be understood as cued, time-compressed readout of the sequences of places ahead of the animal. Such "sweeps" provide information to other brain regions of what is to come. Consider that the place field (210 cm to 270 cm) can be divided into seven subregions, a, b, c, d, e, f, and g. The rat enters the field from the left at position a, which cues the ordered prediction of upcoming positions (Battaglia et al., 2004), thereby producing a "sweep." The "true" place field of this cell is the position g; the firing of this cell at position a represents a prediction that g is ahead of the animal. By the time of the second theta cycle, the rat has moved to b, so b is the cue and produces the sweep b, c, d, e, f, g, h in the second theta cycle. One can see from this example why the g cell fires with earlier theta phase on each successive theta cycle. According to one hypothesis, the successive activation of cells during a sweep results from simple chaining of excitation between the ensembles (Burgess and O'Keefe, 2011; Jensen and Lisman, 1996; Tsodyks et al., 1996). The results shown are combined data from many passages through the place field; the linkage of phase to position is even more exact when individual passages are analyzed (Schmidt et al., 2009), perhaps because theta can occur during different modes (see below) having different phase-position relationships.

(B) Many CA1 place cells fire at a preferred phase of gamma cycles ("in" and "out" refer to the firing probability as a function of gamma phase when the rat is respectively inside or outside of the place field; the line with no error bars is the LFP filtered in the gamma band). Reproduced with permission of Society for Neuroscience, from Senior et al. (2008); permission conveyed through Copyright Clearance Center, Inc. Because the analysis averages over different functional modes (Colgin et al., 2009; de Almeida et al., 2012; Gupta et al., 2012; Senior et al., 2008) and gamma generators (Belluscio et al., 2012), the actual modulation by gamma phase may be stronger than shown.)

Information is represented by an ensemble of cells rather than by single cells. Can a cell assembly that codes for a particular place be observed, and do the cells fire together in a gamma cycle, as postulated? Testing this prediction is difficult because the "true" place field (~3 cm; see Figure 2 caption) is only a small fraction of the environment. Therefore, finding two place cells that code for the same position is difficult, but has been achieved (Dragoi and Buzsáki, 2006). It can be seen in Figure 4 that two cells (red, green) have nearly identical place fields (Figure 4A). The key observation is that these cells tend to fire in the same gamma cycle, as indicated by the fact that the cross-correlation of spiking in the two cells (Figure 4B, red) has a peak that is very near 0 ms and a half-width of ± 15 ms. These results and similar data in (Shapiro and Ferbinteanu, 2006; Skaggs et al., 1996) provide the few glimpses available of actual cell assemblies.

The evidence we have described makes a strong case that the theta-gamma code is used to format multi-item messages about spatial information. We now turn to the question of whether this coding scheme can be linked to cognitive processes and to actual information transfer.




Figure 3. Cells that Fire in Sequence during a Theta Cycle Form a Sweep

Three examples are shown. (Top) Top panels show track. Current position of rat is marked by arrowhead, and it is at this position that the sweep shown below occurs. Different cells active during the sweep have place field centers marked by the colored dots. (Bottom) The spikes of different cells during a theta cycle (start to end). Cells are positioned on the y axis according to the position of their place field centers. In these examples, activity sweeps represent positions along the path ahead of the animal. From Gupta et al. (2012).

Changes in Theta, Gamma, and Their Interactions during Tasks

Although the existence of brain oscillations has been known for many years, the idea that these oscillations provide a mechanistic framework for memory processes is relatively recent and has been controversial. One strategy has been to ask the simple question of whether oscillations are changed during memory processes. Another, more telling approach has been to test whether the magnitude of observed change predicts the accuracy of subsequent memory performance. Initial studies focused on individual types of oscillation; more recent studies have examined the role of theta-gamma coupling. Below, we briefly review these studies—first those on long-term memory and then those on working memory.

Long-Term Memory

Gamma. Studies of long-term memory have focused on the hippocampus. It was found that gamma power and spike-gamma coherence in the monkey hippocampus were higher during successful encoding (Jutras et al., 2009). Similar correlations have been found in humans, both in the hippocampus (Sederberg et al., 2007) and cortical regions (Osipova et al., 2006; Sederberg et al., 2007).

Theta. In rats, hippocampal theta increases during locomotion or attention (Vanderwolf, 1969) and is necessary for memory function (Winson, 1978). In humans, the theta power preceding the stimulus predicted subsequent memory (Fell et al., 2011; Guderian et al., 2009). Using a somewhat different strategy, Rutishauser et al. (2010) showed that successful memory formation was predicted by how well spike timing was phase coupled to theta oscillations. Recent work suggests that, in humans, “slow theta” (3–4 Hz) is predictive of correct recall (Lega et al., 2012; Watrous et al., 2013) and corresponds functionally to the 7 Hz theta of rats.

Theta-Gamma Coupling. Several signal processing tools have been developed to quantify theta-gamma coupling (more generally termed cross-frequency coupling) (Canolty et al., 2006; Cohen, 2008; Kramer et al., 2008; Onslow et al., 2011; Penny et al., 2008; Tort et al., 2010; Young and Eggermont, 2009). These

measure the relationship between the phase of the theta oscillations and the envelope of the gamma power. Thus, high values of coupling indicate that gamma amplitude is a strong function of theta phase. Theta/gamma coupling has

been shown to be functionally important for long-term memory processes (Tort et al., 2009). In this study, rats learned to associate contexts with the location of subsequent food reward (Figure 5A). As learning progressed, there was an increase in cross-frequency coupling (Figure 5B). Moreover, the strength of coupling predicted the probability of correct choice (Figure 5C). These and related results (Shirvalkar et al., 2010) suggest that theta-gamma coupling in the rat hippocampus enables the recall of stored information. A recent study on humans points to a similar conclusion: long-term memory encoding was associated with increased cross-frequency coupling between frontal theta and posterior gamma oscillations (Fries et al., 2012).

Working Memory

In a typical working memory experiment, subjects are presented with a list of several items. This is followed by a delay period (usually <10 s) during which no information is presented. Subjects are then shown a test item and must make a response based on the properties of the information stored in their working memory.

Gamma. In humans, MEG studies and intracranial recordings have reported an increase in gamma power during the delay period of working memory tasks. Importantly, this increase varies with the number of items being held in memory (Howard et al., 2003; Roux et al., 2012; van Vugt et al., 2010). An increase in spike-field coherence during the delay period has also been observed (Pesaran et al., 2002).

Theta. Several studies in humans have reported sustained theta band cortical activity during the delay period (Gevins et al., 1997; Jensen and Tesche, 2002; Raghavachari et al., 2001; Scheeringa et al., 2009), pointing to a role for theta in working memory. One objection to this conclusion is that single-unit recording of persistent firing during a working memory task did not reveal any obvious theta rhythmicity (Funahashi et al., 1991). However, rhythmicity can be difficult to detect by analysis of spikes alone and is more easily detected by determining whether spikes are phase locked to the oscillations in the local field potential (Wang, 2010). Consistent with this, experiments in monkeys have shown that persistent firing during




Figure 4. Cells that Code for the Same Position (Part of a Cell Assembly) and Then Fire in Synchrony (within a Gamma Cycle)
(A) Firing rate as a function of distance on a linear track for five place cells in CA1 (different colors). Green and red cells have peak firing rates at nearly the same position.
(B) The cross-correlation between green and red cells (in A) has a peak very near 0 ms (red line), indicating that the cells tend to fire in the same gamma cycle. This is true even though the particular gamma subcycle changes during the theta-phase precession. Green and blue cells (in A) have different place field centers and tend to fire with a temporal offset that peaks at 40 ms (cyan line) (i.e., in different gamma cycles). From [Dragoi and Buzsáki \(2006\)](#).

a working memory task is phase locked to low-frequency (theta/delta) oscillations in the local field potential, both in extrastriate cortex ([Lee et al., 2005](#); [Liebe et al., 2012](#)) and in prefrontal cortex ([Siegel et al., 2009](#)).

Theta-Gamma Coupling. Theta-gamma coupling during working memory has been demonstrated in humans both in cortex ([Canolty et al., 2006](#); [Lee et al., 2005](#)) and in the hippocampus ([Maris et al., 2011](#)). Gamma power in the hippocampus is modulated by the phase of theta oscillations during working memory retention, and the strength of this cross-frequency coupling predicts individual working memory performance ([Axmacher et al., 2010](#)). Importantly, the particular cortical regions demonstrating cross-frequency coupling depend on the nature of the information being held in working memory. The spatial distribution of gamma band power in cortex can be used to predict whether working memory is maintaining an indoor or outdoor scene ([Fuentemilla et al., 2010](#)), and the gamma activity with this predictive capability is phase locked to the theta activity. In another study, gamma power at certain sites (primarily in occipital cortex) was shown to depend on the particular letter being viewed, and gamma was found to be phase locked to theta ([Jacobs and Kahana, 2009](#)).

The concept of the theta-gamma code ([Lisman and Idiart, 1995](#)) was originally proposed as a way of keeping multiple items active in working memory and there have been several related implementations of this idea ([Jensen and Lisman, 1998](#); [Koene and Hasselmo, 2007](#); [Lisman and Idiart, 1995](#)). Support for this class of models has come from the analysis of grid cells in the entorhinal cortex ([de Almeida et al., 2012](#)), a region that provides input to the hippocampus and has been previously implicated in working memory ([Gaffan and Murray, 1992](#); [Ranganath and D'Esposito, 2001](#); [Stern et al., 2001](#); [Suzuki et al., 1997](#); [Young et al., 1997](#)). It was found that the entorhinal cortex has a working

memory mode in which grid cells represent the recent past (i.e., positions behind the animal). Consistent with the model of [Figure 1B](#), cells representing different positions fired in different gamma subcycles of the theta cycle.

Another way of asking whether the theta-gamma code underlies working memory is to relate the oscillations to the psychophysically measured properties of working memory. A classic result ([Miller, 1956](#)) is that working memory has a capacity limit (span) of 7 ± 2 (see [Cowan \[2001\]](#) for a slightly lower value). The number of gamma cycles within a theta cycle may be what sets the capacity limitation for working memory ([Lisman and Idiart, 1995](#)). Initial efforts to test this concept sought to use the theta-gamma framework to quantitatively account for response time properties of the Sternberg task (i.e., time to respond to whether a given test item was on a short list presented several seconds before). The linear dependence of response time on the number of items in working memory suggested that the list was serially and exhaustively scanned at a rate of 20–30 ms per memory item ([Sternberg, 1966](#)), a time that approximately equals the duration of a gamma cycle. These and other quantitative results of the Sternberg task can be accounted for by models based on the theta-gamma code assuming either that theta phase is reset by stimuli or that theta frequency decreases with memory load ([Jensen and Lisman, 1998](#)). Experiments provide evidence for both effects ([Axmacher et al., 2010](#); [Moran et al., 2010](#); [Mormann et al., 2005](#); [Rizzuto et al., 2006](#)). Recent work sought to determine whether properties of theta and gamma oscillations in individuals could explain their memory span. The ratio of theta to gamma (i.e., the maximum number of gamma cycles within a theta cycle) was found to correlate with span ([Kamiński et al., 2011](#)). However, the determinations of oscillation frequencies were very noise sensitive, raising doubts about the conclusion. Rigorous testing of this relationship will require resolution of the controversy about which brain regions are responsible for short-term memory maintenance and better methods for noninvasive measurement of the oscillatory frequencies at those locations.

Interfering with Oscillations

The experiments described in the previous section make the strong case that theta and gamma oscillations are modified during memory tasks and that these modifications are predictive of performance. However, such experiments are only correlational; a stronger confirmation would require showing that disruption of the oscillations interferes with memory. Experiments of this kind are beginning to be possible because of increased understanding of the mechanisms that underlie oscillations and new optogenetic methods for perturbing the oscillations.

Gamma oscillations are generated by a feedback loop in which firing of pyramidal cells excites fast-spiking interneurons, which then rapidly inhibit the entire population of pyramidal cells (reviewed in [Buzsáki and Wang, 2012](#)). Pyramidal cell firing is re-initiated after the GABAergic inhibition decays, thereby generating the next gamma cycle. Genetic methods have been used to block the excitation of fast-spiking interneurons by reducing their AMPAR-mediated input or by preventing firing using halorhodopsin. Both methods produce a decrease in gamma amplitude and result in memory deficits ([Fuchs et al., 2007](#); [Sohal et al., 2009](#)).




Figure 5. Theta-Gamma Cross-Frequency Coupling in CA3 Predicts Behavior

(A) Exposure to empty context A or B (light gray/dark gray) cues the rat about which cup (blue/yellow) when subsequently presented will contain reward. (B) While exploring context, cross-frequency coupling occurs (left) and can vary in the degree to which theta phase modulates gamma power (right). (C) Theta-gamma modulation occurs during recall of the association between the context and the type of rewarded cup; this modulation is predictive of whether the rat correctly goes to the rewarded cup. From Tort et al. (2009), used with permission.

There is substantial information about the processes that generate theta oscillations in the hippocampus. One important influence is the external input from the medial septal nucleus, a structure that shows theta rhythmicity (King et al., 1998; Zhang et al., 2011). This nucleus contains cholinergic and GABAergic neurons that impose a rhythmic drive onto the hippocampus. However, theta oscillations can also be generated by intrinsic mechanisms, as demonstrated by experiments on the isolated rat hippocampus (Goutagny et al., 2009) and on hippocampal slices (Fellous and Sejnowski, 2000; Fischer et al., 2002; Gloveli et al., 2005). The result of extrinsic and intrinsic influences is a theta frequency inhibition of pyramidal cells, which at its maximum strongly inhibits pyramidal cell firing and reduces the amplitude of gamma oscillations.

Shirvankar et al. (2010) have shown that theta-gamma coupling increases during memory recall. They also found that inhibiting the medial septum with muscimol reduced theta-gamma coupling and interfered with recall. Remarkably stimulating the medial septum at theta frequency partially restored recall. Further evidence for the importance of theta-phase coding has come from experiments examining the effects of the cannabinoid receptor agonist CP5590 (Robbe et al., 2006). This agent strongly disrupted rats' performance on a hippocampal-dependent delayed spatial alternation task. Recordings showed that hippocampal place fields were normal. However, the theta-phase precession normally seen in these cells was completely absent, suggesting that phase-coded information is necessary for this task.

Role of Theta and Gamma Oscillations in Communication between Regions

A strong possibility is that the communication between regions is affected by the coherence of the oscillations in the sender and

receiver networks (Buzsáki, 2010). Coherence occurs when the phase difference in a given frequency band is consistent over time. Coherence varies between 0 and 1, and high values are realistic. For instance, theta coherence can be 0.95 between different parts of the hippocampus (Penley et al., 2012), while theta coherence is in the range of 0.5 between hippocampus and interacting regions in the PFC, amygdala and striatum (Seidenbecher et al., 2003; Sirota et al., 2008; van der Meer and Redish, 2011). The theta coherence between two structures can be elevated in specific phases of a task (Benchenane et al., 2010; Kim et al., 2011; Young and Shapiro, 2011). Importantly, it has been found that during periods of decision, theta coherence between the hippocampus and striatum could be >0.8 and that the magnitude of coherence was predictive of learning (DeCoteau et al., 2007).

As we have argued, the importance of the theta rhythm is to provide a way of ordering multipart messages, an ordering that is exemplified by the phase precession. Phase precession is found both in the structures that provide input to the hippocampus (e.g., the entorhinal cortex and subiculum) (Hafting et al., 2008; Kim et al., 2012; Mizuseki et al., 2009) and in structures to which the hippocampus projects (e.g., the PFC and striatum). Notably, firing in the rat mPFC shows phase precession coupled to that seen in the hippocampus (Jones and Wilson, 2005). Similarly, theta-phase precession can be seen in the striatum (Figure 6; Jones and Wilson, 2005; van der Meer and Redish, 2011). The strong coherence between the hippocampus and its targets and the existence of phase coding in target structures strongly suggest that theta oscillations organize communication between the hippocampus and distant brain regions.

Theta-phase coding is also implicated in communication that does not involve the hippocampus. Theta-phase synchronization occurs between V4 and PFC and is predictive of task




Figure 6. Phase Precession, an Indicator of Theta-Gamma Coupling, Is Evident in the Hippocampus (HC) (Top) but Is Also Evident in a Hippocampal Target Area, the Ventral Striatum (VS) (Bottom)

The data are taken as the rat begins its approach to the reward site. Reproduced with permission of Society for Neuroscience, from van der Meer and Redish (2011); permission conveyed through Copyright Clearance Center, Inc.

performance (Liebe et al., 2012). Although volume conduction confounds are often difficult to assess in human EEG and MEG studies, there are several encouraging findings indicating theta synchronization between frontal and posterior regions in working memory and error-monitoring tasks (Brzezicka et al., 2011; Cavanagh et al., 2009; Cohen and Cavanagh, 2011; Palva et al., 2010; Sarthain et al., 1998; Sauseng et al., 2005; Schack et al., 2005).

The findings summarized above make the strong case that theta oscillations are important for long-range communication and that a signature of such communication is a high level of coherence, as originally proposed by (von Stein and Sarthain, 2000). How this high level of coherence is achieved remains unclear. Within the hippocampus, high theta coherence occurs because the entire structure is driven by a theta generator in the medial septal nucleus of the basal forebrain. A nearby structure, the nucleus basalis, innervates cortex and is a good candidate mechanism for synchronizing cortical theta (Alonso et al., 1996; Lee et al., 2005). However, bidirectional interactions between cortex and thalamus (da Silva et al., 1973; Hughes and Crunelli, 2005; Lopes da Silva et al., 1980; Lorincz et al., 2009; Poulet et al., 2012; Saalman and Kastner, 2011) and intrinsic mechanisms may also be involved (Alonso et al., 1996; Blatow et al., 2003; Flint and Connors, 1996; Jones, 2004; Raghavachari et al., 2006).

There is now substantial evidence suggesting that gamma synchronization between regions can also change during a task. Gamma coherence between monkey parietal and prefrontal areas has been shown to increase from 0.1 to 0.18 during an attention task (Gregoriou et al., 2009). Colgin et al. (2009) found alternating modes in which CA1 became coherent either with entorhinal cortex (through the fast gamma characteristic of the entorhinal region) or with CA3 (through the slower gamma characteristic of CA3). A recent study by Bosman and colleagues provides compelling evidence that gamma coherence reflects the selectivity of attention-mediated communication (Bosman et al., 2012). Two regions in V1 were studied that converged onto V4. When attention was turned to one V1 region,

the coherence of this region with V4 was increased from 0.02 to 0.12 (the other V1 region was not affected). Granger analysis indicated that this change was due to the influence of V1 on V4. Changes in coherence have been correlated with memory performance. A study analyzing data from depth electrodes in epileptic patients showed that gamma synchronization between rhinal cortex and hippocampus predicted memory formation (Fell et al., 2001). In a rat study, gamma band synchronization between CA3 and CA1 reflected performance in a spatial memory task of the behaving rat (Montgomery and Buzsáki, 2007).

An interesting possibility is that changes in gamma coherence may actually control the routing of information (Bressler, 1995; Fries, 2005; Siegel et al., 2012; Varela et al., 2001). This mechanism has been termed the communication through coherence (CTC) hypothesis (Fries, 2005). The general idea is that there are cycles of excitability in oscillatory networks; inputs will be most effective if they arrive at peaks of excitability. Thus, a mechanism that made gamma oscillations in two regions synchronous might selectively route information from one region to the other. However, several difficulties with this hypothesis must be noted. First, the measured levels of long-range gamma coherence are generally very low (0.1–0.2), so any matching of input with the local phase of gamma will be weak (it remains possible that coherence is high but is made low by signal-to-noise problems). Computational studies suggest that strong coherence is required for selective routing (Akam and Kullmann, 2012). Second, there is no indication of an external driver that can impose coherent gamma oscillations in two communicating regions; it is thus thought that coherence develops because of entrainment or resonance mechanisms, processes that develop over many gamma cycles. This process is thus too slow to explain the routing changes that occur during changes in covert attention, which occur in a time period of about a single gamma cycle (Buschman and Miller, 2009; Treisman and Gelade, 1980). These difficulties pose a challenge to the CTC hypothesis; perhaps, to the contrary, increases in gamma coherence are not the cause of selective routing but rather the consequence of some other routing mechanism (see also Rolls et al., 2012). It is unclear what the alternative mechanism might be. One possibility is suggested by recent work showing that the thalamus (pulvinar) can coordinate coherent activity in the alpha bands between parts of visual cortex and produce the routing changes that are required for selective attention (Saalman et al., 2012).

In summary, the presence of information coded by theta phase in the striatum and PFC, together with the strong coherence of theta between the hippocampus and these regions, point to the importance of theta phase coding in long-range communication between brain regions. The role of gamma coherence in long range communication is less clear. In the next section we discuss various postulated roles of gamma. This is followed by a summary section in which we attempt to bring the available information together.

What Does Gamma Contribute to the Theta-Phase Code?

Clocking

Gamma is sometimes referred to as the brain's clock. According to this view, a given gamma cycle would occur at a predictable

time that was a multiple of the gamma period. This view now seems unlikely. Gamma frequency in visual cortex changes with the contrast of visual stimuli (Ray and Maunsell, 2010). Moreover, the gamma period even varies from cycle to cycle (Burns et al., 2011; Henrie and Shapley, 2005; Ray and Maunsell, 2010). This variation occurs in a predictable way: the amplitude of gamma during one cycle (presumably reflecting the number of active cells) determines the duration of the next gamma cycle (Atallah and Scanziani, 2009). The variation of gamma frequency with intensity implies that two parts of the same object having different intensity cannot be synchronized in the same gamma cycles (Ray and Maunsell, 2010), as required if gamma has a binding role in perception (Engel and Singer, 2001; but see Nikolić et al., 2013). In the model of Figure 1B, there is also a concept of binding (cells that represent the same item fire in the same gamma cycle), but it is not assumed that different cells represent different components of the item.

Synchronization and Interitem Separation

Firing occurs during only part of the gamma cycle, a synchronization that allows downstream networks to use coincidence mechanisms to detect ensembles (König et al., 1996). Less recognized is the importance of the pauses that separate the firing that occurs in sequential gamma cycles. As noted earlier, information formatting is a critical aspect of coding that allows downstream “receiver” networks to interpret the coded message (Buzsáki, 2010). In Figure 7, we compare two schemes for organizing the multi-item message in the “sender” network. In one scheme, the sender rapidly switches from a state representing one item to a state representing the next item, i.e., there is no interitem separation. In the other scheme, the sender clusters spikes so that they fire in about one-third of a gamma cycle, thereby creating inter-item separation. Now consider the fact that, for a cell to process input, there must be a window over which it integrates input. Experimentally, this window is shorter than a gamma cycle (Losonczy and Magee, 2006; Pouille and Scanziani, 2001). Without interitem separation, receiver cells will integrate inputs that are part of different items. These cells may therefore detect conjunctions that do not correspond to those in either of the items present. Such false positives are avoided if the pauses generated by gamma frequency inhibition are present.

Selecting Which Cells Fire

A second critical role for gamma is in the *selection* of which cells fire, thereby forming the representation that is active during a gamma cycle. According to one major theory (van Vreeswijk and Sompolinsky, 1996), cells receive large excitation that is balanced by a comparable inhibition; what determines whether a cell fires is stochastic deviations from this balance. However, this theory is not applicable to networks with oscillating inhibition, as discussed by (Isaacson and Scanziani, 2011). An alternative theory (“E%-max”) posits that, in networks with dynamic inhibition, a network process rather than a single-cell process determines which cells fire (de Almeida et al., 2009). According to this theory, the critical step is a “search” for the most excitable cells in the network that occurs as inhibition decays during a gamma cycle. The most excited cells in the network will reach threshold first and trigger global feedback inhibition of less-excited cells (Figure 8). However, this inhibition occurs with




Figure 7. Role of the Firing Pause between Items

Firing as a function of time for two sequential items in schemes with or without a firing pause between items. (Top) An ensemble representing one item (green) followed by rapid state change to an ensemble representing another item (red). (Bottom) Firing for the first item is clustered in one-third of a gamma cycle followed by a pause. Firing for the second item then occurs in the first part of the second gamma cycle, followed by another pause. Downstream networks detect patterns by integrating inputs that arrive within the coincidence detection window (yellow bar). If green is represented by cells 1, 2, 3, red by cells 4, 5, 6, and blue by 1, 2, 6, then the message detected in the state-change case is green, blue, red. In contrast, when firing is clustered, the message is correctly read as green, red.

a delay of a few milliseconds, allowing somewhat less-excited cells to fire before feedback inhibition arrives. Thus, gamma allows the firing of not only the most excited cell, but also somewhat less-excited cells. As a result of this process, the most excited cell (i.e., best tuned to the stimulus) will fire first in a gamma cycle followed by cells that are slightly less well tuned. This prediction has been recently confirmed: it was found that orientation tuning in V1 is highest early in a gamma cycle but becomes slightly lower later in the cycle (Womelsdorf et al., 2012).

Summary

There is now little doubt that multi-item information is formatted by a theta-gamma code in the hippocampus: different spatial locations are represented at different theta phases, and firing is clustered into discrete periods by the gamma rhythm. These oscillations and their interactions are altered during both long-term and working memory processes, as would be expected if different neural operations make different use of these oscillations. That said, we lack a clear understanding of why different tasks (or no task) result in the observed quantitative changes. A critical test of whether the theta-gamma code has an important role in memory is to determine whether inhibiting the oscillations interferes with memory. The available results suggest that this is the case, but more precise experiments are needed. Recently, optogenetic methods have been used to show that odor recognition can be disrupted by selectively interfering with information processes at particular phases of the sniff cycle (Smear et al., 2011). If the hypothesis we are proposing is correct, disrupting information at a particular theta phase should affect information represented at that phase, but not information represented at other phases.

Theta is critical for the transmission of multi-item messages because it provides a phase reference that signifies the onset




Figure 8. How Inhibition during Gamma Oscillations Determines Which Cells Fire

Interneurons fired during a previous gamma cycle (left) and generated global inhibition in pyramidal cells. Inhibition decays with the time constant of the IPSP (τ ; this equals the membrane time constant) (left). The voltage in three pyramidal cells having different tonic excitation is shown (red/green/black). As inhibition decays, the most excited cell (red) reaches threshold. This triggers an interneuron spike and feedback inhibition, which arrives at all pyramidal cells with delay, d . During this delay, the cell (green) with excitation that is close to that of the most excited cell will also fire.

of the message. This phase reference must be shared by sender and receiver; the high observed theta coherence between communicating regions appears to satisfy this requirement. The role of gamma is to define an item in a multi-item message. Gamma contributes to this in three ways: (1) it helps to form the message by allowing only the most excited cells to fire, (2) it synchronizes spikes (clustered spiking can be effectively detected in downstream regions), and (3) it creates pauses between items that prevent errors in decoding the message. The communication of the multipart messages to downstream networks may be aided by coherence in the gamma band, but this is probably not required. We suspect that the small increases in gamma coherence that occur during communication are probably a *result* of effective communication rather than the cause. Because gamma cycles are not of the same duration, detection methods based on exact clocking are not plausible. Thus, although phase-dependent detectors (Jensen, 2001) can be used to detect early versus late items, detection of the information in a specific gamma subcycle does not appear possible. However, many useful functions do not require exact clocking. For instance, according to one model (Fukai, 1999), the sequence of actions to be executed is sent from the hippocampus to the striatum by a theta-gamma code; the striatum stores this sequence and then executes the actions in order, using other information to orchestrate the exact timing of each action. Another useful operation would be the recall of a sequence that contained a salient element such as reward. The detection of this element could be important to downstream networks even if the exact position of that element in the recalled sequence was uncertain. Finally, the entire recalled sequence may be processed (chunked) to represent a higher-level item. Network models that perform such chunking depend on the

ordering of items rather than on exact timing (H. Sanders, B. Kolterman, D. Rinberg, A. Koulakov, and J. Lisman, 2012, Soc. Neurosci., abstract).

Outstanding Questions

How Is the Coherence of Hippocampus, Cortex, and Basal Ganglia Produced?

As described above, when the hippocampus communicates with target regions, the theta in the two regions becomes high. Virtually nothing is known about how this coherence is produced. It seems likely that the basal forebrain and/or the thalamus may be involved, but this remains to be investigated.

What Mechanism Underlies Changes in Cross-Frequency Coupling?

The method for studying cross-frequency coupling are strictly operational and thus do not give insight into the underlying mechanism. For instance, a change in cross-frequency coupling could be explained either by a change in the number of gamma cycles per theta cycle (the duty cycle) or by a change in gamma amplitude with a constant number of gamma cycles per theta cycle. New analytical methods may be helpful in providing a clearer view of what is happening during the observed changes in theta-gamma coupling.

Is There Also a Superfast Gamma-Phase Code?

According to our hypothesis, a representation is formed by all of the cells that fire in a gamma cycle, regardless of their gamma phase. Thus, differences in gamma phase (amounting to only milliseconds) are thought to be unimportant and would be averaged over by the time window of integration in receiver networks. Consistent with the unimportance of gamma phase, position reconstruction methods did not show any advantage to using small theta phase differences that correspond to fractions of a gamma cycle (Harris et al., 2003; Jensen and Lisman, 2000). As noted above, V1 cells with millisecond differences in gamma phase have slightly different orientation tuning. We suspect that such millisecond differences are not important because they are integrated by downstream regions. In memory circuits, an important computation, pattern completion, can be performed during a gamma cycle (de Almeida et al., 2007); dealing with multiple input items within a gamma cycle would be disruptive of this processing. Still, the possibility that the brain uses a superfast code based on millisecond differences in gamma phase must be considered (Fries et al., 2007; Nikolić et al., 2013). Indeed, millisecond differences are important in specialized auditory circuits (Wagner et al., 2005; Yang et al., 2008). Furthermore, possible support for a gamma-phase code comes from single-unit recordings during multi-item working memory in monkeys (Siegel et al., 2009). It was found that neurons representing different memories fired maximally at a different gamma phases, consistent with a superfast gamma-phase code. However, given the analysis method used, it cannot be excluded that cells firing with a different gamma phase were actually firing in different gamma cycles (i.e., with a different low-frequency phase). Indeed, this would be consistent with results showing that hippocampal neurons that fire with different theta phase also fire with different gamma phase (Senior et al., 2008). Further work will thus be needed to clarify this important issue.




Figure 9. Theta Power Is Gated Up at Some Cortical Sites and Down at Others during the Period of Working Memory

(A) With intracranial recordings, large-amplitude theta can be observed in single trials of the Sternberg task. First black bar is the time of cue presentation; second black bar indicates presentation of the item to be remembered; at the red bar, the test item is presented and the subject indicates (at the blue bar) whether it is the same as the remembered item. In other trials, more items were presented.

(B) Recordings from three sites where task (white bar) had little effect on theta power (top), where it produced an increase in theta power (middle), or where it produced a decrease in theta power (bottom). Color code at right indicates theta power. Reproduced with permission of Society for Neuroscience, from Raghavachari et al. (2001); permission conveyed through Copyright Clearance Center, Inc.

What Is the Signature of Engaged Neocortex (Role of Alpha and Theta)?

The properties of theta in the neocortex are much less understood than those in the hippocampus. In the hippocampus, increases in theta power are associated with engagement, but whether this is also true in neocortex is unclear. As shown in Figure 9, there are some sites where theta increases and others where it decreases during a working memory task. Which one is the engaged region? A further critical question is whether oscillations in the theta frequency range should be considered fundamentally different from those in the alpha range? In the hippocampus, these frequencies are lumped together as “theta,” but is this also appropriate for cortex? In the following paragraphs, we discuss these questions.

A classic observation is that alpha power in occipital cortex is high when the visual system is not engaged (Adrian and Matthews, 1934). A cue indicating that attention must be turned to one hemifield leads to a sharp drop in alpha power in the engaged (contralateral) cortex (Worden et al., 2000). In contrast, alpha power in ipsilateral visual cortex may actually increase. This increase is associated with suppression of information that is irrelevant to the task (Thut et al., 2006) and with improved performance (Händel et al., 2011). Further evidence that high alpha power is associated with inhibition is that firing rates,

phosphene detection, and the BOLD signal are all reduced (Haegens et al., 2011; Ritter et al., 2009; Romei et al., 2008). One characteristic of alpha is that neural activity is limited to only about half of the cycle (Bollimunta et al., 2008; Bollimunta et al., 2011; Lakatos et al., 2005). This low-duty cycle (for comparison, see high-duty cycle of firing in Figure 2) probably accounts for why perception is dependent on alpha phase (Busch et al., 2009; Dugué et al., 2011; Mathewson et al., 2009). The suppression of alpha power has also been observed in

other sensory and motor regions when they become engaged (Fontanini and Katz, 2005; Hari and Salmelin, 1997; Pfurtscheller et al., 1997). Importantly, the suppression is even specific to the particular motor subregion that is engaged (Pfurtscheller et al., 1997; see also Miller et al., 2009). Thus, in cortex, high alpha power is an inhibited state, and low alpha power appears to be an indicator of engagement.

Do these “alpha rules” also apply to the only slightly slower cortical oscillations in the theta range? It is important to recall that the frequency bands corresponding to alpha and theta were assigned arbitrarily without consideration of function. Consider Figure 9. If we use hippocampal theta as a model, the elevated theta state would be the engaged state. But based on the “alpha rules,” the lowered theta state would be the engaged state. In favor of the latter, work combining EEG and fMRI has demonstrated that the BOLD signal is high when theta power is low (Michels et al., 2010; Scheeringa et al., 2009). Another perspective on this issue comes from consideration of gamma activity. Gamma is present during the high alpha power state and is phase locked to alpha. As alpha power is decreased, gamma power is increased (Spaak et al., 2012). Importantly, when alpha power falls during task engagement, gamma and its modulation by low-frequency oscillations remain (Sauseng et al., 2009). Indeed, the importance of this cross-frequency

coupling was demonstrated by the fact that the coupling varied with memory load up to saturation and that this saturating load correlated with memory span.

Putting these experimental observations together, we suggest that a signature of engaged cortex is a reduction in the power of alpha/theta (perhaps with a lowering of frequency), modulation of gamma amplitude by the remaining alpha/theta oscillations, and an increase of the gamma duty cycle. Thus, the engaged sites in Figure 9 may be the ones at which alpha/theta decreases, but cross-frequency coupling is nevertheless important. Consistent with this view, cross-frequency coupling between gamma and alpha/theta correlates with the BOLD signal better than other available EEG variables (Wang et al., 2012).

Are There Single Theta Cycle Messages?

There are several reports showing brief periods of gamma activity (gamma bursts) in cortex and striatum (Berke, 2009; Edwards et al., 2005; Freeman, 2003; Howe et al., 2011; Jacobs and Kahana, 2009; Sirota et al., 2008; Yang et al., 2012). The duration of these “gamma bursts” is 100–200 ms. Gamma bursts of similar duration underlie sharp waves in the hippocampal CA3 region (Carr et al., 2012). It is curious that this duration is similar to that of a theta cycle. One could thus argue that, even though continuous theta oscillations are not present in these states, a single theta cycle (and associated gamma) is what is occurring. Indeed, during fast behavioral reactions, there may be time for only a single theta cycle. It will thus be of interest to determine whether perturbations that affect the frequency of continuous theta also affect the duration of gamma bursts.

Is There a Role of Theta-Gamma in Sensory Processing?

Early work in the olfactory cortex provided evidence that theta-gamma interactions are involved in cortical sensory processing (Woolley and Timiras, 1965), and there has recently been renewed interest in this possibility. Recordings from awake, behaving rodents have been made from mitral cells, the output neurons of the olfactory bulb. It was found that a given odor produced brief bursts at a particular phase in the sniff (theta) cycle; different cells had different preferred phases (Shusterman et al., 2011). Firing in the bulb is also phase locked to gamma phase (H. Sanders, B. Kolterman, D. Rinberg, A. Koulakov, and J. Lisman, 2012, Soc. Neurosci., abstract). These results and related work in insects (Laurent et al., 2001) suggest that the signature of an odor is a discrete sequence of ensembles organized by theta-gamma oscillations.

In the auditory system, a model has been proposed in which ongoing syllables and words are segmented by an oscillatory cortical decoder based on nested theta and gamma oscillations (Ghitza, 2011; Giraud and Poeppel, 2012; Peelle and Davis, 2012). This process may involve yet slower oscillations tied to larger lexical units, a hypothesis that relates to an important study (Lakatos et al., 2005) showing that theta oscillations can be nested within slower delta frequency oscillations. This and other recent findings (Fujisawa and Buzsáki, 2011) raise the possibility that a hierarchy of oscillations that include gamma, theta, and delta oscillations organizes sensory and memory functions.

Interest in oscillations in the visual system has been stimulated by the finding that the phase of a stimulus with respect to ongoing cortical alpha/theta oscillations is predictive of the perception threshold, at least for attended stimuli (Busch et al.,

2009; Mathewson et al., 2009). This finding was corroborated by a TMS/EEG study in which the perception of phosphenes was modulated by the phase of both frontal and posterior rhythms in the theta and alpha band (Dugué et al., 2011). Furthermore, the phase of oscillations in the theta-alpha band has been implicated in saccade onset (Drewes and VanRullen, 2011; Hamm et al., 2012). Given that covert attention can move very rapidly (Buschman and Miller, 2009), one possibility is that a theta cycle may be subdivided by faster oscillations, with each of these subcycles representing a different component of the visual scene (Miconi and Vanrullen, 2010).

Conclusion

Neurophysiological studies have provided evidence that a theta-gamma code is used to organize information in the hippocampus and to transmit it to targets in the PFC and striatum. Cognitive studies have shown a strong linkage between theta-gamma oscillations and successfully recalled memories. This body of work lays a foundation for understanding the general problem of how multi-item messages are sent between brain regions, including sensory and possibly motor areas. There is now little doubt that brain oscillations will have an important role in these processes, but major questions remain. The ongoing integration of neurophysiological and cognitive approaches is likely to provide answers to these questions.

ACKNOWLEDGMENTS

The authors gratefully acknowledge The Netherlands Organization for Scientific Research (NWO) VICI grant number: 453-09-002, NIH grants awarded to J.E.L.—5R01MH086518 from the National Institute Of Mental Health and 1R01DA027807 from the National Institute On Drug Abuse. The authors thank Michael Kahana, John Maunsell, Don Katz, Nikolai Axmacher, Dan Pollen, and Honi Sanders for comments on the manuscript.

REFERENCES

- Adrian, E.D., and Matthews, B.H.C. (1934). The Berger rhythm: potential changes from the occipital lobes in man. *Brain* 57, 355–385.
- Akam, T.E., and Kullmann, D.M. (2012). Efficient “communication through coherence” requires oscillations structured to minimize interference between signals. *PLoS Comput. Biol.* 8, e1002760.
- Alonso, A., Khateb, A., Fort, P., Jones, B.E., and Mühlethaler, M. (1996). Differential oscillatory properties of cholinergic and noncholinergic nucleus basalis neurons in guinea pig brain slice. *Eur. J. Neurosci.* 8, 169–182.
- Atallah, B.V., and Scanziani, M. (2009). Instantaneous modulation of gamma oscillation frequency by balancing excitation with inhibition. *Neuron* 62, 566–577.
- Axmacher, N., Henseler, M.M., Jensen, O., Weinreich, I., Elger, C.E., and Fell, J. (2010). Cross-frequency coupling supports multi-item working memory in the human hippocampus. *Proc. Natl. Acad. Sci. USA* 107, 3228–3233.
- Battaglia, F.P., Sutherland, G.R., and McNaughton, B.L. (2004). Local sensory cues and place cell directionality: additional evidence of prospective coding in the hippocampus. *J. Neurosci.* 24, 4541–4550.
- Belluscio, M.A., Mizuseki, K., Schmidt, R., Kempter, R., and Buzsáki, G. (2012). Cross-frequency phase-phase coupling between θ and γ oscillations in the hippocampus. *J. Neurosci.* 32, 423–435.
- Benchenane, K., Peyrache, A., Khamassi, M., Tierney, P.L., Gioanni, Y., Battaglia, F.P., and Wiener, S.I. (2010). Coherent theta oscillations and reorganization of spike timing in the hippocampal- prefrontal network upon learning. *Neuron* 66, 921–936.

- Berke, J.D. (2009). Fast oscillations in cortical-striatal networks switch frequency following rewarding events and stimulant drugs. *Eur. J. Neurosci.* 30, 848–859.
- Blatow, M., Rozov, A., Katona, I., Hormuzdi, S.G., Meyer, A.H., Whittington, M.A., Caputi, A., and Monyer, H. (2003). A novel network of multipolar bursting interneurons generates theta frequency oscillations in neocortex. *Neuron* 38, 805–817.
- Bollimunta, A., Chen, Y., Schroeder, C.E., and Ding, M. (2008). Neuronal mechanisms of cortical alpha oscillations in awake-behaving macaques. *J. Neurosci.* 28, 9976–9988.
- Bollimunta, A., Mo, J., Schroeder, C.E., and Ding, M. (2011). Neuronal mechanisms and attentional modulation of corticostriatal α oscillations. *J. Neurosci.* 31, 4935–4943.
- Bosman, C.A., Schoffelen, J.M., Brunet, N., Oostenveld, R., Bastos, A.M., Womelsdorf, T., Rubehn, B., Stieglitz, T., De Weerd, P., and Fries, P. (2012). Attentional stimulus selection through selective synchronization between monkey visual areas. *Neuron* 75, 875–888.
- Bragin, A., Jandó, G., Nádasdy, Z., Hetke, J., Wise, K., and Buzsáki, G. (1995). Gamma (40–100 Hz) oscillation in the hippocampus of the behaving rat. *J. Neurosci.* 15, 47–60.
- Bressler, S.L. (1995). Large-scale cortical networks and cognition. *Brain Res. Brain Res. Rev.* 20, 288–304.
- Brzezicka, A., Kamiński, M., Kamiński, J., and Blinowska, K. (2011). Information transfer during a transitive reasoning task. *Brain Topogr.* 24, 1–8.
- Burgess, N., and O'Keefe, J. (2011). Models of place and grid cell firing and theta rhythmicity. *Curr. Opin. Neurobiol.* 21, 734–744.
- Burns, S.P., Xing, D., and Shapley, R.M. (2011). Is gamma-band activity in the local field potential of V1 cortex a “clock” or filtered noise? *J. Neurosci.* 31, 9658–9664.
- Busch, N.A., Dubois, J., and VanRullen, R. (2009). The phase of ongoing EEG oscillations predicts visual perception. *J. Neurosci.* 29, 7869–7876.
- Buschman, T.J., and Miller, E.K. (2009). Serial, covert shifts of attention during visual search are reflected by the frontal eye fields and correlated with population oscillations. *Neuron* 63, 386–396.
- Buzsáki, G. (2010). Neural syntax: cell assemblies, synapses, and readers. *Neuron* 68, 362–385.
- Buzsáki, G., and Wang, X.J. (2012). Mechanisms of gamma oscillations. *Annu. Rev. Neurosci.* 35, 203–225.
- Buzsáki, G., Anastassiou, C.A., and Koch, C. (2012). The origin of extracellular fields and currents—EEG, ECoG, LFP and spikes. *Nat. Rev. Neurosci.* 13, 407–420.
- Canolty, R.T., Edwards, E., Dalal, S.S., Soltani, M., Nagarajan, S.S., Kirsch, H.E., Berger, M.S., Barbaro, N.M., and Knight, R.T. (2006). High gamma power is phase-locked to theta oscillations in human neocortex. *Science* 313, 1626–1628.
- Carr, M.F., Karlsson, M.P., and Frank, L.M. (2012). Transient slow gamma synchrony underlies hippocampal memory replay. *Neuron* 75, 700–713.
- Cavanagh, J.F., Cohen, M.X., and Allen, J.J. (2009). Prelude to and resolution of an error: EEG phase synchrony reveals cognitive control dynamics during action monitoring. *J. Neurosci.* 29, 98–105.
- Cohen, M.X. (2008). Assessing transient cross-frequency coupling in EEG data. *J. Neurosci. Methods* 168, 494–499.
- Cohen, M.X., and Cavanagh, J.F. (2011). Single-trial regression elucidates the role of prefrontal theta oscillations in response conflict. *Front. Psychol.* 2. <http://dx.doi.org/10.3389/fpsyg.2011.00030>.
- Colgin, L.L., Denninger, T., Fyhn, M., Hafting, T., Bonnevie, T., Jensen, O., Moser, M.B., and Moser, E.I. (2009). Frequency of gamma oscillations routes flow of information in the hippocampus. *Nature* 462, 353–357.
- Cowan, N. (2001). The magical number 4 in short-term memory: a reconsideration of mental storage capacity. *Behav. Brain Sci.* 24, 87–114, discussion 114–185.
- da Silva, F.H., van Lierop, T.H., Schrijer, C.F., and van Leeuwen, W.S. (1973). Organization of thalamic and cortical alpha rhythms: spectra and coherences. *Electroencephalogr. Clin. Neurophysiol.* 35, 627–639.
- de Almeida, L., Idiart, M., and Lisman, J.E. (2007). Memory retrieval time and memory capacity of the CA3 network: role of gamma frequency oscillations. *Learn. Mem.* 14, 795–806.
- de Almeida, L., Idiart, M., and Lisman, J.E. (2009). A second function of gamma frequency oscillations: an E%-max winner-take-all mechanism selects which cells fire. *J. Neurosci.* 29, 7497–7503.
- de Almeida, L., Idiart, M., Villavicencio, A., and Lisman, J. (2012). Alternating predictive and short-term memory modes of entorhinal grid cells. *Hippocampus* 22, 1647–1651.
- DeCoteau, W.E., Thorn, C., Gibson, D.J., Courtemanche, R., Mitra, P., Kubota, Y., and Graybiel, A.M. (2007). Learning-related coordination of striatal and hippocampal theta rhythms during acquisition of a procedural maze task. *Proc. Natl. Acad. Sci. USA* 104, 5644–5649.
- Demiralp, T., Bayraktaroglu, Z., Lenz, D., Junge, S., Busch, N.A., Maess, B., Ergen, M., and Herrmann, C.S. (2007). Gamma amplitudes are coupled to theta phase in human EEG during visual perception. *Int. J. Psychophysiol.* 64, 24–30.
- Dragoi, G., and Buzsáki, G. (2006). Temporal encoding of place sequences by hippocampal cell assemblies. *Neuron* 50, 145–157.
- Drewes, J., and VanRullen, R. (2011). This is the rhythm of your eyes: the phase of ongoing electroencephalogram oscillations modulates saccadic reaction time. *J. Neurosci.* 31, 4698–4708.
- Dugué, L., Marque, P., and VanRullen, R. (2011). The phase of ongoing oscillations mediates the causal relation between brain excitation and visual perception. *J. Neurosci.* 31, 11889–11893.
- Edwards, E., Soltani, M., Deouell, L.Y., Berger, M.S., and Knight, R.T. (2005). High gamma activity in response to deviant auditory stimuli recorded directly from human cortex. *J. Neurophysiol.* 94, 4269–4280.
- Engel, A.K., and Singer, W. (2001). Temporal binding and the neural correlates of sensory awareness. *Trends Cogn. Sci.* 5, 16–25.
- Fell, J., Klaver, P., Lehnertz, K., Grunwald, T., Schaller, C., Elger, C.E., and Fernández, G. (2001). Human memory formation is accompanied by rhinal-hippocampal coupling and decoupling. *Nat. Neurosci.* 4, 1259–1264.
- Fell, J., Ludowig, E., Staresina, B.P., Wagner, T., Kranz, T., Elger, C.E., and Axmacher, N. (2011). Medial temporal theta/alpha power enhancement precedes successful memory encoding: evidence based on intracranial EEG. *J. Neurosci.* 31, 5392–5397.
- Fellous, J.M., and Sejnowski, T.J. (2000). Cholinergic induction of oscillations in the hippocampal slice in the slow (0.5–2 Hz), theta (5–12 Hz), and gamma (35–70 Hz) bands. *Hippocampus* 10, 187–197.
- Fischer, Y., Wittner, L., Freund, T.F., and Gähwiler, B.H. (2002). Simultaneous activation of gamma and theta network oscillations in rat hippocampal slice cultures. *J. Physiol.* 539, 857–868.
- Flint, A.C., and Connors, B.W. (1996). Two types of network oscillations in neocortex mediated by distinct glutamate receptor subtypes and neuronal populations. *J. Neurophysiol.* 75, 951–957.
- Fontanini, A., and Katz, D.B. (2005). 7 to 12 Hz activity in rat gustatory cortex reflects disengagement from a fluid self-administration task. *J. Neurophysiol.* 93, 2832–2840.
- Freeman, W.J. (2003). The wave packet: an action potential for the 21st century. *J. Integr. Neurosci.* 2, 3–30.
- Fries, P. (2005). A mechanism for cognitive dynamics: neuronal communication through neuronal coherence. *Trends Cogn. Sci.* 9, 474–480.
- Fries, P., Nikolić, D., and Singer, W. (2007). The gamma cycle. *Trends Neurosci.* 30, 309–316.
- Friese, U., Köster, M., Hassler, U., Martens, U., Trujillo-Barreto, N., and Gruber, T. (2012). Successful memory encoding is associated with increased cross-frequency coupling between frontal theta and posterior gamma oscillations in human scalp-recorded EEG. *Neuroimage* 66C, 642–647.

- Fuchs, E.C., Zivkovic, A.R., Cunningham, M.O., Middleton, S., Lebeau, F.E., Bannerman, D.M., Rozov, A., Whittington, M.A., Traub, R.D., Rawlins, J.N., and Monyer, H. (2007). Recruitment of parvalbumin-positive interneurons determines hippocampal function and associated behavior. *Neuron* 53, 591–604.
- Fuentemilla, L., Penny, W.D., Cashdollar, N., Bunzeck, N., and Düzel, E. (2010). Theta-coupled periodic replay in working memory. *Curr. Biol.* 20, 606–612.
- Fujisawa, S., and Buzsáki, G. (2011). A 4 Hz oscillation adaptively synchronizes prefrontal, VTA, and hippocampal activities. *Neuron* 72, 153–165.
- Fukai, T. (1999). Sequence generation in arbitrary temporal patterns from theta-nested gamma oscillations: a model of the basal ganglia-thalamo-cortical loops. *Neural Netw.* 12, 975–987.
- Funahashi, S., Bruce, C.J., and Goldman-Rakic, P.S. (1991). Neuronal activity related to saccadic eye movements in the monkey's dorsolateral prefrontal cortex. *J. Neurophysiol.* 65, 1464–1483.
- Gaffan, D., and Murray, E.A. (1992). Monkeys (*Macaca fascicularis*) with rhinal cortex ablations succeed in object discrimination learning despite 24-hr inter-trial intervals and fail at matching to sample despite double sample presentations. *Behav. Neurosci.* 106, 30–38.
- Gevins, A., Smith, M.E., McEvoy, L., and Yu, D. (1997). High-resolution EEG mapping of cortical activation related to working memory: effects of task difficulty, type of processing, and practice. *Cereb. Cortex* 7, 374–385.
- Ghitza, O. (2011). Linking speech perception and neurophysiology: speech decoding guided by cascaded oscillators locked to the input rhythm. *Front. Psychol.* 2. <http://dx.doi.org/10.3389/fpsyg.2011.00130>.
- Giraud, A.L., and Poeppel, D. (2012). Cortical oscillations and speech processing: emerging computational principles and operations. *Nat. Neurosci.* 15, 511–517.
- Gloveli, T., Dugladze, T., Rotstein, H.G., Traub, R.D., Monyer, H., Heinemann, U., Whittington, M.A., and Kopell, N.J. (2005). Orthogonal arrangement of rhythm-generating microcircuits in the hippocampus. *Proc. Natl. Acad. Sci. USA* 102, 13295–13300.
- Goutagny, R., Jackson, J., and Williams, S. (2009). Self-generated theta oscillations in the hippocampus. *Nat. Neurosci.* 12, 1491–1493.
- Gregoriou, G.G., Gotts, S.J., Zhou, H., and Desimone, R. (2009). High-frequency, long-range coupling between prefrontal and visual cortex during attention. *Science* 324, 1207–1210.
- Guderian, S., Schott, B.H., Richardson-Klavehn, A., and Düzel, E. (2009). Medial temporal theta state before an event predicts episodic encoding success in humans. *Proc. Natl. Acad. Sci. USA* 106, 5365–5370.
- Gupta, A.S., van der Meer, M.A., Touretzky, D.S., and Redish, A.D. (2012). Segmentation of spatial experience by hippocampal θ sequences. *Nat. Neurosci.* 15, 1032–1039.
- Haegens, S., Nacher, V., Luna, R., Romo, R., and Jensen, O. (2011). α -Oscillations in the monkey sensorimotor network influence discrimination performance by rhythmical inhibition of neuronal spiking. *Proc. Natl. Acad. Sci. USA* 108, 19377–19382.
- Hafting, T., Fyhn, M., Bonnevie, T., Moser, M.B., and Moser, E.I. (2008). Hippocampus-independent phase precession in entorhinal grid cells. *Nature* 453, 1248–1252.
- Hamm, J.P., Dyckman, K.A., McDowell, J.E., and Clementz, B.A. (2012). Precue fronto-occipital alpha phase and distributed cortical oscillations predict failures of cognitive control. *J. Neurosci.* 32, 7034–7041.
- Händel, B.F., Haarmeier, T., and Jensen, O. (2011). Alpha oscillations correlate with the successful inhibition of unattended stimuli. *J. Cogn. Neurosci.* 23, 2494–2502.
- Hari, R., and Salmelin, R. (1997). Human cortical oscillations: a neuromagnetic view through the skull. *Trends Neurosci.* 20, 44–49.
- Harris, K.D., Csicsvari, J., Hirase, H., Dragoi, G., and Buzsáki, G. (2003). Organization of cell assemblies in the hippocampus. *Nature* 424, 552–556.
- Henrie, J.A., and Shapley, R. (2005). LFP power spectra in V1 cortex: the graded effect of stimulus contrast. *J. Neurophysiol.* 94, 479–490.
- Howard, M.W., Rizzuto, D.S., Caplan, J.B., Madsen, J.R., Lisman, J., Aschenbrenner-Scheibe, R., Schulze-Bonhage, A., and Kahana, M.J. (2003). Gamma oscillations correlate with working memory load in humans. *Cereb. Cortex* 13, 1369–1374.
- Howe, M.W., Atallah, H.E., McCool, A., Gibson, D.J., and Graybiel, A.M. (2011). Habit learning is associated with major shifts in frequencies of oscillatory activity and synchronized spike firing in striatum. *Proc. Natl. Acad. Sci. USA* 108, 16801–16806.
- Hughes, S.W., and Crunelli, V. (2005). Thalamic mechanisms of EEG alpha rhythms and their pathological implications. *Neuroscientist* 11, 357–372.
- Isaacson, J.S., and Scanziani, M. (2011). How inhibition shapes cortical activity. *Neuron* 72, 231–243.
- Jacobs, J., and Kahana, M.J. (2009). Neural representations of individual stimuli in humans revealed by gamma-band electrocorticographic activity. *J. Neurosci.* 29, 10203–10214.
- Jensen, O. (2001). Information transfer between rhythmically coupled networks: reading the hippocampal phase code. *Neural Comput.* 13, 2743–2761.
- Jensen, O., and Lisman, J.E. (1996). Hippocampal CA3 region predicts memory sequences: accounting for the phase precession of place cells. *Learn. Mem.* 3, 279–287.
- Jensen, O., and Lisman, J.E. (1998). An oscillatory short-term memory buffer model can account for data on the Sternberg task. *J. Neurosci.* 18, 10688–10699.
- Jensen, O., and Lisman, J.E. (2000). Position reconstruction from an ensemble of hippocampal place cells: contribution of theta phase coding. *J. Neurophysiol.* 83, 2602–2609.
- Jensen, O., and Tesche, C.D. (2002). Frontal theta activity in humans increases with memory load in a working memory task. *Eur. J. Neurosci.* 15, 1395–1399.
- Johnson, A., and Redish, A.D. (2007). Neural ensembles in CA3 transiently encode paths forward of the animal at a decision point. *J. Neurosci.* 27, 12176–12189.
- Jones, B.E. (2004). Activity, modulation and role of basal forebrain cholinergic neurons innervating the cerebral cortex. *Prog. Brain Res.* 145, 157–169.
- Jones, M.W., and Wilson, M.A. (2005). Phase precession of medial prefrontal cortical activity relative to the hippocampal theta rhythm. *Hippocampus* 15, 867–873.
- Jutras, M.J., Fries, P., and Buffalo, E.A. (2009). Gamma-band synchronization in the macaque hippocampus and memory formation. *J. Neurosci.* 29, 12521–12531.
- Kamiński, J., Brzezicka, A., and Wróbel, A. (2011). Short-term memory capacity (7 ± 2) predicted by theta to gamma cycle length ratio. *Neurobiol. Learn. Mem.* 95, 19–23.
- Kim, J., Delcasso, S., and Lee, I. (2011). Neural correlates of object-in-place learning in hippocampus and prefrontal cortex. *J. Neurosci.* 31, 16991–17006.
- Kim, S.M., Ganguli, S., and Frank, L.M. (2012). Spatial information outflow from the hippocampal circuit: distributed spatial coding and phase precession in the subiculum. *J. Neurosci.* 32, 11539–11558.
- King, C., Recce, M., and O'Keefe, J. (1998). The rhythmicity of cells of the medial septum/diagonal band of Broca in the awake freely moving rat: relationships with behaviour and hippocampal theta. *Eur. J. Neurosci.* 10, 464–477.
- Koene, R.A., and Hasselmo, M.E. (2007). First-in-first-out item replacement in a model of short-term memory based on persistent spiking. *Cereb. Cortex* 17, 1766–1781.
- König, P., Engel, A.K., and Singer, W. (1996). Integrator or coincidence detector? The role of the cortical neuron revisited. *Trends Neurosci.* 19, 130–137.
- Kramer, M.A., Tort, A.B., and Kopell, N.J. (2008). Sharp edge artifacts and spurious coupling in EEG frequency comodulation measures. *J. Neurosci. Methods* 170, 352–357.

- Lakatos, P., Shah, A.S., Knuth, K.H., Ulbert, I., Karmos, G., and Schroeder, C.E. (2005). An oscillatory hierarchy controlling neuronal excitability and stimulus processing in the auditory cortex. *J. Neurophysiol.* *94*, 1904–1911.
- Laurent, G., Stopfer, M., Friedrich, R.W., Rabinovich, M.I., Volkovskii, A., and Abarbanel, H.D. (2001). Odor encoding as an active, dynamical process: experiments, computation, and theory. *Annu. Rev. Neurosci.* *24*, 263–297.
- Lee, H., Simpson, G.V., Logothetis, N.K., and Rainer, G. (2005). Phase locking of single neuron activity to theta oscillations during working memory in monkey extrastriate visual cortex. *Neuron* *45*, 147–156.
- Lega, B.C., Jacobs, J., and Kahana, M. (2012). Human hippocampal theta oscillations and the formation of episodic memories. *Hippocampus* *22*, 748–761.
- Lenck-Santini, P.P., Fenton, A.A., and Muller, R.U. (2008). Discharge properties of hippocampal neurons during performance of a jump avoidance task. *J. Neurosci.* *28*, 6773–6786.
- Liebe, S., Hoerzer, G.M., Logothetis, N.K., and Rainer, G. (2012). Theta coupling between V4 and prefrontal cortex predicts visual short-term memory performance. *Nat. Neurosci.* *15*, 456–462.
- Lisman, J., and Buzsáki, G. (2008). A neural coding scheme formed by the combined function of gamma and theta oscillations. *Schizophr. Bull.* *34*, 974–980.
- Lisman, J.E., and Idiart, M.A. (1995). Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science* *267*, 1512–1515.
- Llinás, R., and Ribary, U. (1993). Coherent 40-Hz oscillation characterizes dream state in humans. *Proc. Natl. Acad. Sci. USA* *90*, 2078–2081.
- Lopes da Silva, F.H., Vos, J.E., Mooibroek, J., and Van Rotterdam, A. (1980). Relative contributions of intracortical and thalamo-cortical processes in the generation of alpha rhythms, revealed by partial coherence analysis. *Electroencephalogr. Clin. Neurophysiol.* *50*, 449–456.
- Lorincz, M.L., Kékesi, K.A., Juhász, G., Crunelli, V., and Hughes, S.W. (2009). Temporal framing of thalamic relay-mode firing by phasic inhibition during the alpha rhythm. *Neuron* *63*, 683–696.
- Losonczy, A., and Magee, J.C. (2006). Integrative properties of radial oblique dendrites in hippocampal CA1 pyramidal neurons. *Neuron* *50*, 291–307.
- Maris, E., van Vugt, M., and Kahana, M. (2011). Spatially distributed patterns of oscillatory coupling between high-frequency amplitudes and low-frequency phases in human iEEG. *Neuroimage* *54*, 836–850.
- Mathewson, K.E., Gratton, G., Fabiani, M., Beck, D.M., and Ro, T. (2009). To see or not to see: prestimulus alpha phase predicts visual awareness. *J. Neurosci.* *29*, 2725–2732.
- Mehta, M.R., Lee, A.K., and Wilson, M.A. (2002). Role of experience and oscillations in transforming a rate code into a temporal code. *Nature* *417*, 741–746.
- Michels, L., Bucher, K., Lüchinger, R., Klaver, P., Martin, E., Jeanmonod, D., and Brandeis, D. (2010). Simultaneous EEG-fMRI during a working memory task: modulations in low and high frequency bands. *PLoS ONE* *5*, e10298.
- Miconi, T., and Vanrullen, R. (2010). The gamma slideshow: object-based perceptual cycles in a model of the visual cortex. *Front. Hum. Neurosci.* *4*. <http://dx.doi.org/10.3389/fnhum.2010.00205>.
- Miller, G.A. (1956). The magical number seven plus or minus two: some limits on our capacity for processing information. *Psychol. Rev.* *63*, 81–97.
- Miller, K.J., Zanos, S., Fetz, E.E., den Nijs, M., and Ojemann, J.G. (2009). Decoupling the cortical power spectrum reveals real-time representation of individual finger movements in humans. *J. Neurosci.* *29*, 3132–3137.
- Mizuseki, K., Sirota, A., Pastalkova, E., and Buzsáki, G. (2009). Theta oscillations provide temporal windows for local circuit computation in the entorhinal-hippocampal loop. *Neuron* *64*, 267–280.
- Montgomery, S.M., and Buzsáki, G. (2007). Gamma oscillations dynamically couple hippocampal CA3 and CA1 regions during memory task performance. *Proc. Natl. Acad. Sci. USA* *104*, 14495–14500.
- Moran, R.J., Campo, P., Maestu, F., Reilly, R.B., Dolan, R.J., and Strange, B.A. (2010). Peak frequency in the theta and alpha bands correlates with human working memory capacity. *Front. Hum. Neurosci.* *4*. <http://dx.doi.org/10.3389/fnhum.2010.00200>.
- Mormann, F., Fell, J., Axmacher, N., Weber, B., Lehnertz, K., Elger, C.E., and Fernández, G. (2005). Phase/amplitude reset and theta-gamma interaction in the human medial temporal lobe during a continuous word recognition memory task. *Hippocampus* *15*, 890–900.
- Nikolić, D., Fries, P., and Singer, W. (2013). Gamma oscillations: precise temporal coordination without a metronome. *Trends Cogn. Sci.* *17*, 54–55.
- O'Keefe, J., and Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat. *Brain Res.* *34*, 171–175.
- O'Keefe, J., and Recce, M.L. (1993). Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* *3*, 317–330.
- Onslow, A.C., Bogacz, R., and Jones, M.W. (2011). Quantifying phase-amplitude coupling in neuronal network oscillations. *Prog. Biophys. Mol. Biol.* *105*, 49–57.
- Osipova, D., Takashima, A., Oostenveld, R., Fernández, G., Maris, E., and Jensen, O. (2006). Theta and gamma oscillations predict encoding and retrieval of declarative memory. *J. Neurosci.* *26*, 7523–7531.
- Palva, S., Monto, S., and Palva, J.M. (2010). Graph properties of synchronized cortical networks during visual working memory maintenance. *Neuroimage* *49*, 3257–3268.
- Pastalkova, E., Itskov, V., Amarasingham, A., and Buzsáki, G. (2008). Internally generated cell assembly sequences in the rat hippocampus. *Science* *321*, 1322–1327.
- Peelle, J.E., and Davis, M.H. (2012). Neural oscillations carry speech rhythm through to comprehension. *Front. Psychol.* *3*. <http://dx.doi.org/10.3389/fpsyg.2012.00320>.
- Penley, S.C., Hinman, J.R., Sabolek, H.R., Escabi, M.A., Markus, E.J., and Chrobak, J.J. (2012). Theta and gamma coherence across the septotemporal axis during distinct behavioral states. *Hippocampus* *22*, 1164–1175.
- Penny, W.D., Duzel, E., Miller, K.J., and Ojemann, J.G. (2008). Testing for nested oscillation. *J. Neurosci. Methods* *174*, 50–61.
- Penttonen, M., Kamondi, A., Acsády, L., and Buzsáki, G. (1998). Gamma frequency oscillation in the hippocampus of the rat: intracellular analysis in vivo. *Eur. J. Neurosci.* *10*, 718–728.
- Pesaran, B., Pezaris, J.S., Sahani, M., Mitra, P.P., and Andersen, R.A. (2002). Temporal structure in neuronal activity during working memory in macaque parietal cortex. *Nat. Neurosci.* *5*, 805–811.
- Pfurtscheller, G., Neuper, C., Andrew, C., and Edlinger, G. (1997). Foot and hand area mu rhythms. *Int. J. Psychophysiol.* *26*, 121–135.
- Pouille, F., and Scanziani, M. (2001). Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science* *293*, 1159–1163.
- Poulet, J.F., Fernandez, L.M., Crochet, S., and Petersen, C.C. (2012). Thalamic control of cortical states. *Nat. Neurosci.* *15*, 370–372.
- Raghavachari, S., Kahana, M.J., Rizzuto, D.S., Caplan, J.B., Kirschen, M.P., Bourgeois, B., Madsen, J.R., and Lisman, J.E. (2001). Gating of human theta oscillations by a working memory task. *J. Neurosci.* *21*, 3175–3183.
- Raghavachari, S., Lisman, J.E., Tully, M., Madsen, J.R., Bromfield, E.B., and Kahana, M.J. (2006). Theta oscillations in human cortex during a working-memory task: evidence for local generators. *J. Neurophysiol.* *95*, 1630–1638.
- Ranganath, C., and D'Esposito, M. (2001). Medial temporal lobe activity associated with active maintenance of novel information. *Neuron* *31*, 865–873.
- Ray, S., and Maunsell, J.H. (2010). Differences in gamma frequencies across visual cortex restrict their possible use in computation. *Neuron* *67*, 885–896.
- Ritter, P., Moosmann, M., and Villringer, A. (2009). Rolandic alpha and beta EEG rhythms' strengths are inversely related to fMRI-BOLD signal in primary somatosensory and motor cortex. *Hum. Brain Mapp.* *30*, 1168–1187.

- Rizzuto, D.S., Madsen, J.R., Bromfield, E.B., Schulze-Bonhage, A., and Kahana, M.J. (2006). Human neocortical oscillations exhibit theta phase differences between encoding and retrieval. *Neuroimage* *31*, 1352–1358.
- Robbe, D., Montgomery, S.M., Thome, A., Rueda-Orozco, P.E., McNaughton, B.L., and Buzsáki, G. (2006). Cannabinoids reveal importance of spike timing coordination in hippocampal function. *Nat. Neurosci.* *9*, 1526–1533.
- Rolls, E.T., Webb, T.J., and Deco, G. (2012). Communication before coherence. *Eur. J. Neurosci.* *36*, 2689–2709.
- Romei, V., Brodbeck, V., Michel, C., Amedi, A., Pascual-Leone, A., and Thut, G. (2008). Spontaneous fluctuations in posterior alpha-band EEG activity reflect variability in excitability of human visual areas. *Cereb. Cortex* *18*, 2010–2018.
- Roux, F., Wibral, M., Mohr, H.M., Singer, W., and Uhlhaas, P.J. (2012). Gamma-band activity in human prefrontal cortex codes for the number of relevant items maintained in working memory. *J. Neurosci.* *32*, 12411–12420.
- Rutishauser, U., Ross, I.B., Mamelak, A.N., and Schuman, E.M. (2010). Human memory strength is predicted by theta-frequency phase-locking of single neurons. *Nature* *464*, 903–907.
- Saalmann, Y.B., and Kastner, S. (2011). Cognitive and perceptual functions of the visual thalamus. *Neuron* *71*, 209–223.
- Saalmann, Y.B., Pinsk, M.A., Wang, L., Li, X., and Kastner, S. (2012). The pulvinar regulates information transmission between cortical areas based on attention demands. *Science* *337*, 753–756.
- Sarnthein, J., Petsche, H., Rappelsberger, P., Shaw, G.L., and von Stein, A. (1998). Synchronization between prefrontal and posterior association cortex during human working memory. *Proc. Natl. Acad. Sci. USA* *95*, 7092–7096.
- Sauseng, P., Klimesch, W., Schabus, M., and Doppelmayr, M. (2005). Frontoparietal EEG coherence in theta and upper alpha reflect central executive functions of working memory. *Int. J. Psychophysiol.* *57*, 97–103.
- Sauseng, P., Klimesch, W., Heise, K.F., Gruber, W.R., Holz, E., Karim, A.A., Glennon, M., Gerloff, C., Birbaumer, N., and Hummel, F.C. (2009). Brain oscillatory substrates of visual short-term memory capacity. *Curr. Biol.* *19*, 1846–1852.
- Schack, B., Klimesch, W., and Sauseng, P. (2005). Phase synchronization between theta and upper alpha oscillations in a working memory task. *Int. J. Psychophysiol.* *57*, 105–114.
- Scheeringa, R., Petersson, K.M., Oostenveld, R., Norris, D.G., Hagoort, P., and Bastiaansen, M.C. (2009). Trial-by-trial coupling between EEG and BOLD identifies networks related to alpha and theta EEG power increases during working memory maintenance. *Neuroimage* *44*, 1224–1238.
- Schmidt, R., Diba, K., Leibold, C., Schmitz, D., Buzsáki, G., and Kempter, R. (2009). Single-trial phase precession in the hippocampus. *J. Neurosci.* *29*, 13232–13241.
- Sederberg, P.B., Schulze-Bonhage, A., Madsen, J.R., Bromfield, E.B., McCarthy, D.C., Brandt, A., Tully, M.S., and Kahana, M.J. (2007). Hippocampal and neocortical gamma oscillations predict memory formation in humans. *Cereb. Cortex* *17*, 1190–1196.
- Seidenbecher, T., Laxmi, T.R., Stork, O., and Pape, H.C. (2003). Amygdalar and hippocampal theta rhythm synchronization during fear memory retrieval. *Science* *301*, 846–850.
- Senior, T.J., Huxter, J.R., Allen, K., O'Neill, J., and Csicsvari, J. (2008). Gamma oscillatory firing reveals distinct populations of pyramidal cells in the CA1 region of the hippocampus. *J. Neurosci.* *28*, 2274–2286.
- Shapiro, M.L., and Ferbinteanu, J. (2006). Relative spike timing in pairs of hippocampal neurons distinguishes the beginning and end of journeys. *Proc. Natl. Acad. Sci. USA* *103*, 4287–4292.
- Shirvalkar, P.R., Rapp, P.R., and Shapiro, M.L. (2010). Bidirectional changes to hippocampal theta-gamma comodulation predict memory for recent spatial episodes. *Proc. Natl. Acad. Sci. USA* *107*, 7054–7059.
- Shusterman, R., Smear, M.C., Koulakov, A.A., and Rinberg, D. (2011). Precise olfactory responses tile the sniff cycle. *Nat. Neurosci.* *14*, 1039–1044.
- Siegel, M., Warden, M.R., and Miller, E.K. (2009). Phase-dependent neuronal coding of objects in short-term memory. *Proc. Natl. Acad. Sci. USA* *106*, 21341–21346.
- Siegel, M., Donner, T.H., and Engel, A.K. (2012). Spectral fingerprints of large-scale neuronal interactions. *Nat. Rev. Neurosci.* *13*, 121–134.
- Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M., and Buzsáki, G. (2008). Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta rhythm. *Neuron* *60*, 683–697.
- Skaggs, W.E., McNaughton, B.L., Wilson, M.A., and Barnes, C.A. (1996). Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* *6*, 149–172.
- Smear, M., Shusterman, R., O'Connor, R., Bozza, T., and Rinberg, D. (2011). Perception of sniff phase in mouse olfaction. *Nature* *479*, 397–400.
- Sohal, V.S., Zhang, F., Yizhar, O., and Deisseroth, K. (2009). Parvalbumin neurons and gamma rhythms enhance cortical circuit performance. *Nature* *459*, 698–702.
- Soltesz, I., and Deschênes, M. (1993). Low- and high-frequency membrane potential oscillations during theta activity in CA1 and CA3 pyramidal neurons of the rat hippocampus under ketamine-xylazine anesthesia. *J. Neurophysiol.* *70*, 97–116.
- Spaak, E., Bonnefond, M., Maier, A., Leopold, D.A., and Jensen, O. (2012). Layer-specific entrainment of gamma-band neural activity by the alpha rhythm in monkey visual cortex. *Curr. Biol.* *22*, 2313–2318.
- Stern, C.E., Sherman, S.J., Kirchhoff, B.A., and Hasselmo, M.E. (2001). Medial temporal and prefrontal contributions to working memory tasks with novel and familiar stimuli. *Hippocampus* *11*, 337–346.
- Sternberg, S. (1966). High-speed scanning in human memory. *Science* *153*, 652–654.
- Suzuki, W.A., Miller, E.K., and Desimone, R. (1997). Object and place memory in the macaque entorhinal cortex. *J. Neurophysiol.* *78*, 1062–1081.
- Thut, G., Nietzel, A., Brandt, S.A., and Pascual-Leone, A. (2006). Alpha-band electroencephalographic activity over occipital cortex indexes visuospatial attention bias and predicts visual target detection. *J. Neurosci.* *26*, 9494–9502.
- Tort, A.B., Komorowski, R.W., Manns, J.R., Kopell, N.J., and Eichenbaum, H. (2009). Theta-gamma coupling increases during the learning of item-context associations. *Proc. Natl. Acad. Sci. USA* *106*, 20942–20947.
- Tort, A.B., Komorowski, R., Eichenbaum, H., and Kopell, N. (2010). Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. *J. Neurophysiol.* *104*, 1195–1210.
- Treisman, A.M., and Gelade, G. (1980). A feature-integration theory of attention. *Cognit. Psychol.* *12*, 97–136.
- Tsodyks, M.V., Skaggs, W.E., Sejnowski, T.J., and McNaughton, B.L. (1996). Population dynamics and theta rhythm phase precession of hippocampal place cell firing: a spiking neuron model. *Hippocampus* *6*, 271–280.
- van der Meer, M.A., and Redish, A.D. (2011). Theta phase precession in rat ventral striatum links place and reward information. *J. Neurosci.* *31*, 2843–2854.
- van Vreeswijk, C., and Sompolinsky, H. (1996). Chaos in neuronal networks with balanced excitatory and inhibitory activity. *Science* *274*, 1724–1726.
- van Vugt, M.K., Schulze-Bonhage, A., Litt, B., Brandt, A., and Kahana, M.J. (2010). Hippocampal gamma oscillations increase with memory load. *J. Neurosci.* *30*, 2694–2699.
- Vanderwolf, C.H. (1969). Hippocampal electrical activity and voluntary movement in the rat. *Electroencephalogr. Clin. Neurophysiol.* *26*, 407–418.
- Varela, F., Lachaux, J.P., Rodriguez, E., and Martinerie, J. (2001). The brainweb: phase synchronization and large-scale integration. *Nat. Rev. Neurosci.* *2*, 229–239.
- von Stein, A., and Sarnthein, J. (2000). Different frequencies for different scales of cortical integration: from local gamma to long range alpha/theta synchronization. *Int. J. Psychophysiol.* *38*, 301–313.

- Voytek, B., Canolty, R.T., Shestyuk, A., Crone, N., Parvizi, J., and Knight, R.T. (2010). Shifts in gamma phase-amplitude coupling frequency from theta to alpha over posterior cortex during visual tasks. *Front. Hum. Neurosci.* *4*. <http://dx.doi.org/10.3389/fnhum.2010.00191>.
- Wagner, H., Brill, S., Kempner, R., and Carr, C.E. (2005). Microsecond precision of phase delay in the auditory system of the barn owl. *J. Neurophysiol.* *94*, 1655–1658.
- Wang, X.J. (2010). Neurophysiological and computational principles of cortical rhythms in cognition. *Physiol. Rev.* *90*, 1195–1268.
- Wang, L., Saalman, Y.B., Pinsk, M.A., Arcaro, M.J., and Kastner, S. (2012). Electrophysiological low-frequency coherence and cross-frequency coupling contribute to BOLD connectivity. *Neuron* *76*, 1010–1020.
- Watrous, A.J., Lee, D.J., Izadi, A., Gurkoff, G.G., Shahlaie, K., and Ekstrom, A.D. (2013). A comparative study of human and rat hippocampal low frequency oscillations during spatial navigation. *Hippocampus*, in press.
- Wilson, M.A., and McNaughton, B.L. (1993). Dynamics of the hippocampal ensemble code for space. *Science* *261*, 1055–1058.
- Winson, J. (1978). Loss of hippocampal theta rhythm results in spatial memory deficit in the rat. *Science* *201*, 160–163.
- Womelsdorf, T., Lima, B., Vinck, M., Oostenveld, R., Singer, W., Neuenschwander, S., and Fries, P. (2012). Orientation selectivity and noise correlation in awake monkey area V1 are modulated by the gamma cycle. *Proc. Natl. Acad. Sci. USA* *109*, 4302–4307.
- Woolley, D.E., and Timiras, P.S. (1965). Prepyriform electrical activity in the rat during high altitude exposure. *Electroencephalogr. Clin. Neurophysiol.* *18*, 680–690.
- Worden, M.S., Foxe, J.J., Wang, N., and Simpson, G.V. (2000). Anticipatory biasing of visuospatial attention indexed by retinotopically specific alpha-band electroencephalography increases over occipital cortex. *J. Neurosci.* *20*, RC63.
- Yang, Y., DeWeese, M.R., Otazu, G.H., and Zador, A.M. (2008). Millisecond-scale differences in neural activity in auditory cortex can drive decisions. *Nat. Neurosci.* *11*, 1262–1263.
- Yang, J.-W., An, S., Sun, J.-J., Reyes-Puerta, V., Kindler, J., Berger, T., Kilb, W., and Luhmann, H.J. (2012). Thalamic network oscillations synchronize ontogenetic columns in the newborn rat barrel cortex. *Cereb. Cortex*. Published online May 16, 2012. <http://dx.doi.org/10.1093/cercor/bhs103>.
- Young, C.K., and Eggermont, J.J. (2009). Coupling of mesoscopic brain oscillations: recent advances in analytical and theoretical perspectives. *Prog. Neurobiol.* *89*, 61–78.
- Young, J.J., and Shapiro, M.L. (2011). Dynamic coding of goal-directed paths by orbital prefrontal cortex. *J. Neurosci.* *31*, 5989–6000.
- Young, B.J., Otto, T., Fox, G.D., and Eichenbaum, H. (1997). Memory representation within the parahippocampal region. *J. Neurosci.* *17*, 5183–5195.
- Zhang, H., Lin, S.C., and Nicolelis, M.A. (2011). A distinctive subpopulation of medial septal slow-firing neurons promote hippocampal activation and theta oscillations. *J. Neurophysiol.* *106*, 2749–2763.